

Amendments to the claims:

Claims 1-28. (Cancelled)

Claim 29. (Previously presented) A method for selecting an efficacious therapy for a subject suffering from a schizophrenia, said method comprising the steps of:

- (a) analyzing the MTHFR nucleic acid in a sample obtained from said subject;
- (b) determining the presence of a heterozygous C/T mutation at position 677 that decreases MTHFR activity or reduces MTHFR thermal stability MTHF in said subject, wherein the presence of said mutation is indicative of the efficacy of a therapy; and
- (c) selecting an efficacious neuroleptic therapy for said subject.

Claim 30 and 31. (Cancelled)

Claim 32. (Previously presented) The method of claim 29, wherein said nucleic acid with said mutation at position 677 is determined to further comprise an additional mutation selected from the group consisting of: G/A mutation at position 167, a G/A mutation at position 482, a C/T mutation a position 559, a C/T mutation at position 692, a C/T mutation at position 764, a G/A mutation at position 792+1, a C/T mutation at position 985, a C/T mutation at position 1015, a C/T mutation at position 1081, an A/C

mutation at position 1298, and a T/C mutation at position 1317, wherein said additional mutation decreases MTHFR activity or reduces MTHFR thermal stability.

Claim 33. (Previously presented) The method of claim 32, wherein said nucleic acid with said mutation at position 677 comprises an A/C mutation at position 1298.

Claim 34. (Previously presented) The method of claim 29, wherein said nucleic acid is determined to comprise at least two MTHFR mutations at a position other than 677 that further decrease MTHFR activity or further reduce MTHFR thermal stability.

Claim 35. (Previously presented) The method of claim 34, wherein said mutations at a position other than 677 comprise at least one of a G/A mutation at position 167, a G/A mutation at position 482, a C/T mutation a position 559, a C/T mutation at position 692, a C/T mutation at position 764, a G/A mutation at position 792+1, a C/T mutation at position 985, a C/T mutation at position 1015, a C/T mutation at position 1081, an A/C mutation at position 1298, or a T/C mutation at position 1317.

Claim 36. (Previously presented) The method of claim 35, wherein said mutations at a position other than 677 comprise an A/C mutation at position 1298.

Claims 37-44. (Cancelled)

Claim 45. (Previously presented) A method for treating a schizophrenia in a subject, said method comprising the steps of:

- (a) analyzing the MTHFR nucleic acid in a sample obtained from said subject;
- (b) determining the presence of a heterozygous C/T mutation at position 677 that decreases MTHFR activity or reduces MTHFR thermal stability in said subject, wherein the presence of said mutation is predictive of the responsiveness of said schizophrenia to a neuroleptic therapy; and
- (c) administering said neuroleptic therapy to said subject.

Claims 46 and 47. (Cancelled)

Claim 48. (Previously presented) The method of claim 45, wherein said nucleic acid with said mutation at position 677 further comprises a G/A mutation at position 167, a G/A mutation at position 482, a C/T mutation a position 559, a C/T mutation at position 692, a C/T mutation at position 764, a G/A mutation at position 792+1, a C/T mutation at position 985, a C/T mutation at position 1015, a C/T mutation at position 1081, an A/C mutation at position 1298, or a T/C mutation at position 1317, wherein said additional mutation decreases MTHFR activity or reduces MTHFR thermal stability.

Claim 49. (Previously presented) The method of claim 48, wherein said nucleic acid with said mutation at position 677 comprises an A/C mutation at position 1298.

Claim 50. (Previously presented) The method of claim 45, wherein said nucleic acid is determined to comprise at least two mutations at a position other than 677 that further decrease MTHFR activity or further reduce MTHFR thermal stability.

Claim 51. (Previously presented) The method of claim 50, wherein said mutations at a position other than 677 comprise at least one of a G/A mutation at position 167, a G/A mutation at position 482, a C/T mutation a position 559, a C/T mutation at position 692, a C/T mutation at position 764, a G/A mutation at position 792+1, a C/T mutation at position 985, a C/T mutation at position 1015, a C/T mutation at position 1081, an A/C mutation at position 1298, or a T/C mutation at position 1317.

Claim 52. (Previously presented) The method of claim 51, wherein said mutations at a position other than 677 comprise an A/C mutation at position 1298.

Claim 53. (Previously presented) The method of claim 29, further comprising the step of determining the presence of at least one MTHFR mutation at a position other than

677 prior to step (c) wherein said mutation decreases MTHFR activity or reduces MTHFR thermal stability.

Claim 54. (Cancelled)

Claim 55. (Previously presented) The method of claim 45, further comprising determining the presence of at least one MTHFR mutation at a position other than 677 prior to step (c), wherein said additional mutation decreases MTHFR activity or reduces MTHFR thermal stability.

Claim 56. (Previously presented) The method of claim 29, wherein said step (c) further comprises selecting a folic acid therapy.

Claim 57. (Previously presented) The method of claim 45, wherein said step (c) further comprises administering a folic acid therapy.

Claims 58-64. (Cancelled)